

Focal dermal hypoplasia: a case report and review of literature

* A. O. Ogunbiyi¹, I. O. Adewole², O. Ogunleye³, J. O. Ogunbiyi⁴,
O. O. Ogunseinde⁵ and A. Baiyeroju-Agbeja⁶

¹Department of Medicine, University College Hospital, Ibadan, ²Department of Obstetrics & Gynaecology, University College Hospital, Ibadan, ³Department of Surgery, University College Hospital, Ibadan, ⁴Department of Pathology, University College Hospital, Ibadan, ⁵Department of Radiology, University College Hospital, Ibadan and ⁶Department of Ophthalmology, University College Hospital, Ibadan, Nigeria.

Summary

A 14 year old girl was seen in the clinic with a huge vulvar wart, and warts on both tonsils. She also had poikiloderma. She had bony abnormalities, which included lobster claw abnormality of the right foot, a right cervical rib, and the right clavicle was lower than the left. There were fine parallel vertical radio-opaque lines in the distal femoral tibial metaphyses bilaterally (osteopathia striata).

A diagnosis of Focal dermal hypoplasia was made.

The huge vulvar wart and the warts on the tonsils were excised. She recovered promptly and she was discharged home.

Keywords: *Focal dermal hypoplasia, Nigerian, Case report*

Résumé

Une fille âgée de 14 ans était vue dans la clinique atteinte d'un défaut énorme dans la vulve, et les défauts sur les deux amygdales. Elle est également atteinte de la poikilodermie. Elle a mal dans l'os y compris une anomalie dans le lobster claw de la jambe droite, une côte cervicale du droite et la clavicle du droite plus basse que celle de gauche. Il y a des lignes radio-opaque fines parallèles verticales bilatéralement dans les métaphyses distale femorale (osteopathia striata).

On a fait un diagnostique de la hypoplasie dermal focal. Le défaut très énorme de la vulve et les défauts sur les amygdales ont été opérés. Elle s'est rétabli rapidement et sortie de l'hôpital.

Introduction

Focal dermal hypoplasia is an uncommon disorder affecting organs derived from the ecto and mesoderm. Its cutaneous features were first described in 1921¹ and the non dermatological components of the disorder were highlighted by Goltz in 1962.² One of the unique features of the disorder is its variability in organ involvement. The skin, eyes, and skeletal tissue are usually involved. One of the components of the syndrome, papillomas may be present at birth or appear at different sites later on in life. The patients may present with life threatening situations e.g. papillomas may develop in the larynx requiring tracheostomy. The papillomas may also mimic infections such as genital warts which may be wrongly attributed to child abuse in children. We present a case of FDH in a 14 years old Nigerian female who presented with papillomas on the tonsils and perianal area. She also had a capillary haemangioma in the right gluteal region. As far as we know haemangioma has not been previously documented in this syndrome and it is the first case reported in a Nigerian.

Case report

A 14 year old Nigerian girl was seen in the Gynaecological department of the University College Hospital Ibadan with a five month history of a vulvar mass which was progressively increasing in size. The mass bled especially when it was traumatized and exuded foul smelling fluid. She denied any history of sexual intercourse in the past. Her mother had noticed that she snored during sleep and her voice had become somewhat muffled in the last two years.

There were no associated systemic symptoms and the patient felt well otherwise. Her past medical history revealed that she had presented to the same hospital at the age of 4 years with a complaint of a right gluteal mass, which had been present since birth but had increased in size at that time. The mass had been excised and the histology report from the notes and review of the slides at this visit showed the mass to be a capillary haemangioma (Fig. 1). There had been no recurrence of the lesion. She was also noticed to have shortening of the right limb at that visit.

Table 1 Skeletal abnormalities in FDH

Short stature
Asymmetry of skull, trunk and extremities
Hypoplastic or absent digits
Syndactyly
Osteoporosis ²⁴
Fibrous dysplasia ²⁴
Cyst in metatarsals, fibula, tibia, maxilla, ileum and ischium
Widening of public symphysis ^{4,25}
Abnormalities of ribs and clavicles
Lobster claw hand and foot
Vertebral abnormalities e.g. scoliosis
Osteopathia striata ^{26,27}
Osteochondroma ⁴ or Giant cell tumours ²⁸



Fig. 1 Shows capillary haemangioma (H & E x 100)

Her mother claimed she had an uneventful pregnancy and delivered her at nine months. She could not remember her birth weight but claimed that she had some raw areas on the skin of the lower limb at birth which had healed within three weeks. She also had a congenital abnormality of the right foot.

The mother denied any history of similar disorder in the family. She did not have any history of miscarriages at any time. She had four other children who were all normal. She also felt her daughter was not mentally retarded, but had started school late because of frequent hospital visits.

Examination of the 14 year old revealed, a young girl with a small stature, weighing 25kg and measuring 1.36 meters in height. Her head circumference was 50cm.

She had a triangular face with large ears (Fig. 2)

Examination of the skin revealed linear streaks of hypo and hyperpigmentation that followed Blaschko's lines. The trunk, limbs,

* Correspondence




Fig. 2 Patient with FDH with hypo and hyperpigmented streaks following Blaschko's lines mainly on the right. There is shortening of the right tibia and fibula.




Fig. 4 Chest radiograph showing cervical ribs (arrows) and left clavicle higher than the right.



Fig. 5 Xray of the right knee joint showing linear sclerotic bands osteopathia striata



Fig. 3 Numerous papillomas on both tonsils which appear to meet in the middle "kissing tonsils"

and part of the face were affected. The pigmentary abnormalities were interspersed with striate atrophic areas and the right side of the body was more affected. (Fig. 2) Erythema was prominent in the hypopigmented area of the lower limbs. She had scarring alopecia in areas affected by the atrophic lesions. The third fingernail was absent and this had been so from birth.

The breasts were normally developed for her size but she had sparse axillary and public hair. She had scoliosis and shortening of the right tibia and fibula by about 6cm. She also had a lobster claw deformity of the right foot.

Eye examination revealed posterior subcapsular cataract.

Oral examination revealed small conical shaped teeth with malocclusion.

She had markedly enlarged tonsils with wart-like growths that appeared to meet in the midline "kissing tonsils" (Fig. 3). The larynx appeared normal.

Examination of her genitals revealed a huge right vulvar growth




Fig. 6 Section of skin biopsy showing absence of rete pegs and dermal appendages with bundles of mature adipocytes within the dermis. (H & E x 40)

extending to the perianal area with a lobulated surface. It had a few ulcers on the surface and exuded blood stained foul smelling fluid. Her hymen was intact. The rectal examination was normal.

Examination of the cardiovascular, respiratory, and central nervous systems appeared normal. Results of investigations are outlined below.

Her haematological work up and electrolytes and urea were within normal limits. X-ray findings were rounded lucencies in the

parietal bones in keeping with congenital parietal foramina. Both frontal sinuses were dysplastic, while the ethmoids were cloudy with the right nasal turbinate engorged, suggesting chronic sinusitis. There was caries of the left second upper incisor and rarefaction of the maxillary bone anteriorly. There was a right cervical rib and a prominent transverse process on the left side, with the left clavicle being higher than the right (Fig. 4). X-ray of the thighs and knees showed linear lucencies in the muscles of the thighs especially on the posterior aspect. There were fine parallel vertical radio-opaque lines in the distal femoral and proximal tibial metaphyses bilaterally, better demonstrated on the right side, osteopathia striata (Fig. 5). There was also loss of parallelism of the articular surfaces of the right knee due to uneven modeling and expansion of the medial notch. Pelvic ultrasound revealed a hypoplastic uterus.

She then had diathermy excision of the vulvar growth and bilateral tonsillectomy. Histology of the tonsils and vulvar mass showed acanthosis, papillomatosis and koilocytosis of the epithelium with focal parakeratosis. Some of the tonsillar crypts were covered by debris and foamy histiocytes. These lesions were thought to be hamatomatous.

Histology of the atrophic skin lesion showed loss of rete pegs with bundles of mature adipocytes within the reticular and papillary dermis associated with atrophy of dermal appendages (Fig. 6). These features were consistent with atrophy that could be associated with Goltz Syndrome. The patient subsequently had an eventful recovery from the surgery and was discharged home.

Discussion

The term 'focal dermal hypoplasia' does not describe the protean manifestations of this disorder that involves various organs in the body but it has come to stay. FDH is a developmental disorder of organs of mesodermal and ectodermal origin. There are about 200 cases in the literature.³ There is strong evidence that it is inherited as an X linked dominant condition.⁴ It has a female preponderance and there is an increased frequency of miscarriage of male fetuses in affected females.^{4,5} However a few cases have been documented in males^{6,7,8} and these have been attributed to half chromatid mutations or autosomal dominant inheritance affecting the germ cells.⁹ Spontaneous cases resulting from mutation do occur as can be assumed to be the case in our patient with no family history of the disorder nor mosaicism of it.

The variability in the expression of FDH within families and involvement of the skin and bone can be explained by the Lyon hypothesis in which random inactivation of the mutant gene results in mosaicism,¹⁰ and the percentage of the active mutant X chromosome present would determine the clinical picture in the individual. The atrophic linear streaks with pigmentary abnormalities following Blaschko's lines on the skin, and the linear streaks on the bone indicate some form of X chromosome mosaicism resulting in clonal proliferation of two functionally different populations of cells during embryogenesis.¹¹

Some authors have confirmed the findings of two different populations of dermal fibroblasts in the affected individual with fibroblasts in the affected skin being abnormal while that in the unaffected skin was normal.^{12,13}

FDH manifests itself in different organs of ectodermal and mesodermal origin but its effect is mainly on the connective tissue of bone and skin.^{3,14} Patients are of small stature, they have a triangular face and large ears for the size of their face. Cutaneous manifestations of FDH in the same individual may be varied. It may be bilateral but asymmetric or solely unilateral. There are reports of patients with FDH without skin involvement¹⁵ but skin involvement is usually regarded as essential for diagnosis.

When hair and nail disorders¹⁶ are excluded, the major cutaneous features could be grouped into five.

1. Congenital aplasia of the skin
2. Multiform patterns of atrophic like lesions
3. Striate lesions.
4. Verrucoid papilloma of skin and mucous membrane (angiofibroma).
5. Lipomatous lesions

Telangiectasia and hyperpigmentation of various designs and amounts may accompany or overlap the multiform atrophic lesions and striated lipomatous lesions.

Congenital aplasia of the skin

This is not very common and the skin is completely absent in the affected site at birth.^{17,18} In some patients there may be some bullous eruptions or inflammatory process on the skin at birth which usually heals up leaving a scar as was reported in our patient.

Multiform pattern of atrophic like depression

This has been described with different names such as atrophy, depressions, scars, congenital linear muscular atrophoderma or focal dermal hypoplasia. Histological features of these lesions depend on the age of the lesion. New lesions may show an increase in dermal blood vessels but the older lesions show a thin or absent dermis, i.e. dermal hypoplasia with mid dermal fat deposits.

Striate lesions

These are linear stripes or streaks which are congenital. They may be inflammatory at the onset and may be confused with incontinentia pigmenti. The histology of Striate lesions usually reveals increased proliferation of blood vessels, increase in ground substances, and infiltration of lymphocytes and histiocytes.

Papillomas

The papillomatous lesions known as angiofibroma or fibrovascular papillomas are an important component of the disorder that may be present at birth or develop as the child gets older. They have been reported in various locations in the body but the commonest sites are the perineum, vulva, and perianal regions.^{3,16,19} The vulvar lesions are usually misdiagnosed as genital warts (condyloma acuminata)^{16,17} as was the case in our patient. The warts on her tonsils had suggested the possibility of oral sex which in the younger patients may insinuate sexual abuse and lead to false accusations. Papillomas have been reported in the larynx, eye lid margins, mouth, pharynx, tonsils, palate, gums, tongue and lips. The fingers and pinna may also have papillomas on them. These lesions may also produce life threatening situations e.g. papillomas in the larynx requiring tracheotomy,²⁰ or those occurring in the oesophagus causing stricture.²¹ Malignant transformation of papillomas has also been reported.²²

Histology of the papillomas show a fibrovascular stalk covered with a layer of acanthotic stratified squamous epithelium resembling epidermis with extensive papillary folds. Hyperkeratosis and parakeratosis may be present. The stalk is composed of loose connective tissue with dilated vessels and an admixture of inflammatory cells.

Lipomatosis

These manifest as skin coloured nodules or nodules slightly darker than the normal skin. Lipomatosis does not occur in all patients and was not present in our patient. The histology of the lesions show lobulated fat approaching the epidermis but separated by a few strands of connective tissue. This results from hypoplasia of the dermis which varies in the different areas. The hair may be sparse especially if the scalp is involved and apocrine gland abnormalities have been reported.²²

Hydrocystomas have also been reported around the eyes.²³ Nail abnormalities include dystrophic nails or total absence of the

nail as was seen in our patient.

Skeletal abnormalities in the patients with FDH have been highlighted in table 1. Of note is the linear streaks in the metaphyses of the long bones osteopathia striata. This has established itself as one of the hallmarks of FDH^{26,27} which was also seen in our patient. It is not present in all cases. The lobster claw deformity affecting the hands or foot are also striking abnormalities of the syndrome. A number of abnormalities of soft tissue have also been reported in a few of these patients. They include urinary tract abnormalities e.g. horse shoe kidney, dilated ureters with inflammation and cystitis. Others include a diaphragmatic hernia and reflux oesophagitis. Some of the patients do mature and have reproductive capabilities but our patient had a hypoplastic uterus.

Excellent reviews on eye signs in FDH have been published^{29,30} and they include colobomas, (especially of the iris), microphthalmia, anophthalmia, hypertelorism, optic atrophy, keratoconu, cortical and subcapsular cataract. Strabismus, nystagmus and disorders of lachrymal gland apparatus have also been documented.

Many patients with FDH have some level of mental deficiency and there are reports of meningomyelocele, hydrocephalus and Arnold Chiari malformation in a patient.³¹ Seizures have not been reported in association with this disorder. Our patient had an associated capillary haemangioma on the right gluteal region which is in keeping with mesodermal abnormalities. As far as we know this has not been previously reported in association with this syndrome.

References

- Ruiz Maldonado R, Carnevale A and Tamayo L et al: Focal dermal hypoplasia. *Clin Genet* 1974; 6: 36 - 35.
- Goltz RW, Paterson WC and Gorlin RJ et al: Focal dermal hypoplasia. *Arch Dermatol* 1962; 86: 708-717.
- Goltz RW: Focal dermal hypoplasia syndrome: an update. *Arch Dermatol* 1992; 18: 812 - 5.
- Holden JD and Akers WA: Goltz syndrome: Focal dermal hypoplasia. *AJDC* 1967; 114: 292 - 300.
- Wettke-schaefer R and Kantner G: X linked dominantly inherited diseases with lethality in hemizygous males. *Human Genet* 1983; 84: 1 - 24.
- Toro-sola MA, Kistenmacher ML, Punnett HH and Di George: Focal dermal hypoplasia syndrome in a male. *Clin Genet* 1970; 325 - 327.
- Feinberg A and Menter MA: Focal dermal hypoplasia (Goltz syndrome) in a male; a case report. *Afr Med J* 1976; 50: 554 - 555.
- Fryns JP, Diondt F, Lindemans L and Van der berghe: Focal dermal hypoplasia (Goltz syndrome) in a male. *Acta Paed* 1978; 31: 37 - 39.
- Temple IK, MacDowall and Baraitser M et al: Focal dermal hypoplasia (Goltz syndrome). *J Med Genet* 1990; 27: 180 - 7.
- Lyon MF: Gene action in X chromosome of the mouse. *Nature* 1961; 190: 372 - 73.
- Wechsler M A, Papa O M, Haberman F and Marion R W: Variable expression in focal dermal hypoplasia: an example of differential X chromosome differentiation.
- Utto J, Bauer E A, Santa Cruz D J, Loewinger R J and Eisen A Z: Focal dermal hypoplasia: Abnormal growth characteristics of skin fibroblasts in culture. *J. Invest dermatol* 1980; 170 - 175.
- Manzi G, Magli A, Pignalosa B and Liquori G: The Goltz syndrome: case report. *Ophthalmologica* 1990; 200: 104 - 6.
- Goltz R W, Hitch J M and Ott J E: Focal dermal hypoplasia syndrome: *Arch dermatol* 1970; 101: 1 - 11.
- Ayne S and Fraser F C: Possible examples of the Goltz syndrome (Focal dermal hypoplasia) without linear areas of skin hypoplasia. *Birth defects* 1982; 18: 59 - 65.
- Howell J B and Freeman R G: Gutaneous defects of focal dermal hypoplasia: an ecto mesodermal dysplasia syndrome. *J Cutan Pathol* 1989; 16: 237 - 258.
- Harken J H and Caplan R M: Aplasia cutis congenita. *Arch Dermatol* 1970; 102: 386 - 389.
- Chikara A J, Anand N K and Saini L L: Aplasia cutis congenita. *Int J. Peadiatr* 1985; 22: 69 - 71.
- Griffiths WAD: Case report. *Proc R Soc Med* 1967; 69: 17 - 18.
- Begovic D and Zergollern L: The Goltz - Gorlin syndrome in a male child. *Lijec vjem.* 1989; 111: 85 - 88.
- Brison R R, Schumen B M, Mills L R and Thigpen S: Multiple squamous papillomas of the oesophagus associated with Goltz syndrome. *Am J Gastroenterol* 1987; 82: 1177 - 79.
- Vakilzadeh F, Happle R, Peters P and Machers E: Focal dermal hypoplasia with apocrine nevi and striation of the back. *Arch Dermatol* 1985; 56: 189 - 195.
- Zala L, Ettin C and Krebs A: Focal dermal hypoplasia with keratoconus, papillomatosis of the oesophagus and hydrocystomas. *Dermatologica* 1975; 150: 176 - 85.
- Cottenot F, Bougeois - Droin C, Wallach D and Guetrot B: Focal dermal hypoplasia with fibrous dysplasia of bone. *Ann Dermatol Venerol* 1979; 106: 167 - 169.
- Beganovic N and Lommen E J: A case of focal dermal hypoplasia (Goltz syndrome) with some new aspects. *Acta Paediatr scand* 1977; 66: 255 - 256.
- Howell J B and Reynolds J: Osteopathia striata - a diagnostic mark of focal dermal hypoplasia. *Trans St. Johns Hosp derm soc.* 1974; 60: 172-182.
- Larreqe Y, Maroteaux P, Michel Y and Faire C: Osteopathia striata-Radiological symptoms of a hypoplastic dermis. *Ann Radiol* 1872; 15: 287-295.
- Joannides T et al: Giant cell tumour of bone in Focal dermal hypoplasia. *Br. J. Radiol* 1983; 56: 684.
- Willet G S: Focal dermal hypoplasia: *Br. J. Ophthalmol* 1974; 58: 620 - 624.
- Thomas J V, Yoshizumi M O, Beyer C K, Craft J L and Albert D M: Ocular manifestations of focal dermal hypoplasia. *Arch Ophthalmol* 1977; 95: 1977 - 2000.
- Almeida L, Anyane-Yeboah K, Grossman M and Rosen T: Meningomyelocele, Arnold-Chiari anomaly and hydrocephalus in focal dermal hypoplasia. *Am J. Genet* 1988; 30: 917 - 923.