

www.ajbrui.net

Afr. J. Biomed. Res. Vol.15 (January 2012); 35- 40

Research Article

Phytochemical Screening and Microbial Inhibitory Activities of *Ficus Capensis*

Adebayo-Tayo B.C and *Odeniyi A.O

Department of Microbiology, University of Ibadan, Ibadan, Nigeria

ABSTRACT: *Ficus* plant components have application in traditional medicine because of the myriad uses they have been subjected to. The ease of application is based on the secondary metabolites this plant contains. The challenges faced by modern medicine especially in the complete cure of microbially-associated diseases through abrupt and unpredictable genetic mutations in the presence of conventional drugs informed the investigation of the microbial inhibitory activities of the stem, root and leaf parts of *F. capensis* against test disease causing microorganisms. The phenolic, alkaloid and tannin phytochemical fractions were highest in *F. capensis* bark extract (180, 165 and 155 µg/ml respectively) followed by that contained in the stem extract (100, 90 and 85 µg/ml respectively). While *Streptococcus faecalis* and *Pseudomonas mirabilis* were resistant to many different antibiotics (87.5%), they were effectively inhibited by all concentrations of ethanolic *F. capensis* extracts. The minimum inhibitory concentration of ethanolic extracts ranged from 25% leaf and stem extract concentration respectively (4mm) against *S. faecalis* and (2mm) against *P.mirabilis*. All test isolates were 100% susceptible to ethanol extract growth inhibition..

Keywords: *Ficus capensis*; phytochemical; inhibitory activities.

INTRODUCTION

Ficus plants belong to the mulberry family, *Moraceae*. *Ficus capensis*, also known as the cape fig and is a native of tropical Africa and the Cape Islands. The plant is a deciduous tree with spreading roots and branches and broad green leaves. The name 'fig' is derived from the hollow, pear-shaped inflorescence called "Syconia". *F. capensis* produces fleshy fruits all year round in a single or branched raceme along the trunk and the main braches. These fruits, though inedible to humans are eaten by many animals.

Many of the world's population are versed in the use of the available plants and herbs in their environment in the treatment of different diseases (Hansch *et al.*, 1990; Shellard, 1987). In Nigeria, and

other parts of the world, the plant along with many others in this family is important in the traditional treatment of many diseases and ailments. The plant extracts have been reported in the treatment of diarrhea, dysentery, sexually transmitted disease causing microorganisms, chest ailments, tuberculosis, leprosy, convulsions, pain, in anaemia and wound (Irvine, 1961; Amos *et al.*, 2001; Ahmadu *et al.*, 2007, Oyeleke *et al.*, 2008; Sandabe and Kwari, 2000; Wakeel *et al.*, 2004) among many others.

The relevance of this plant in traditional medicine is as a result of the secondary metabolites such as phytates, phenols, saponins, tannins, alkaloids, terpenoids and flavonoids which they have been screened to contain. Also referred to as phytochemicals, they are reported to possess inhibitory activities against the growth and disease inducing activities of some pathogenic microorganisms (Hassan, 2005; Oyeleke *et al.*, 2008; Sandabe *et al.*, 2006; Solomon-Wisdom *et al.*, 2011 Stary, 1998; Udobi *et al.*, 2008).

Despite the breakthroughs in disease management and medicine, disease causing microorganisms continually undergo genetic variations which permit them to survive and resist treatment especially with antibiotics. Hence, even in this modern age the search

*Correspondence: (E-mail): busolaodeniya@gmail.com

Received: September 2011; Accepted: November 2011

faecalis, *E. coli* and *S. dysenteriae*. In both cases the inhibition reduced as the concentration of the extract reduced. Water extracts from the leaf and bark demonstrated half of the inhibitory potentials of the ethanol extracts. No inhibition was recorded by either extracts at 0.125g/ml concentration in *P. aeruginosa*, *S. aureus*, *K. pneumoniae* and *S. dysenteriae*. The water extract from the stem was totally without activity against *S. faecalis*, *S. aureus* and *K. aeruginosa*. The minimum inhibitory concentration was recorded from the leaf and stem ethanol fractions against *S. faecalis* and *P. mirabilis*. A concentration of 125µg/ml leaf and stem ethanolic extract respectively was found to effect an inhibition on *S. faecalis* and *P. mirabilis* respectively.

Table 3 shows the response of test isolates to some antibiotics. All isolates were resistant to cotrimoxazole except *S. dysenteriae* which showed an inhibitory zone of 23mm. *K. pneumonia*, *P. mirabilis*, *S. aureus* and *S. faecalis* were all resistant to Tetracycline amoxicillin and nitrosufuranide. *P. mirabilis* and *S. faecalis* had the

greatest resistance to the range of antibiotics tested against being susceptible to only one antibiotic each.

Table 3 shows the sensitivity of the test isolates to some commercial antibiotics. *S. faecalis* was resistant to all antibiotics except nalidixine while all concentrations of the ethanolic extracts had inhibition against it (Table 2). *S. dysenteriae* was also highly resistant to most of the antibiotics and recorded a very high inhibition with the leaf ethanolic extract (16mm). All extracts at 25% to 100% concentration inhibited the bacterium's growth around the well. The highest concentration of the ethanol extracts recorded high inhibition against *K. pneumoniae* while the bacterium proved resistant to 6 antibiotics. A hundred percent concentration of the *F. capensis* leaf ethanol extract recorded an inhibition of 18mm on *E. coli* which was more than the value that was recorded with nitrofurantoin and gentamicin. Inhibition of *P. aeruginosa* by 100% bark ethanol extract (18mm) was higher than that recorded from 5 antibiotics against the same bacterium.




Figure 1: Quantitative determination of the bioactive components of *Ficus capensis* plant parts

Table 2:

Inhibitory activities of *Ficus capensis* extracts on selected isolates

Test Isolates	Extract concentration (%)	Water control	Ethanol control	Zone of inhibition (mm)					
				LEE	SEE	BEE	LWE	SWE	BWE
<i>S. faecalis</i>	100	-	1	13	10	11	8	-	8
	75			8	8	8	2	-	6
	50			8	6	5	-	-	-
	25			4	-	-	-	-	-
<i>P. mirabilis</i>	100	-	1	8	8	5	4	2	5
	75			3	6	2	4	-	3
	50			-	6	-	-	-	-
	25			-	2	-	-	-	-
<i>P. aeruginosa</i>	100	-	2	14	13	18	6	4	5
	75			9	11	6	4	2	-
	50			4	-	-	-	-	-
	25			-	-	-	-	-	-
<i>S. aureus</i>	100	-	1	8	10	12	5	-	2
	75			8	4	3	2	-	-
	50			5	-	3	-	-	-
	25			-	-	-	-	-	-
<i>E. coli</i>	100	-	2	18	15	16	10	8	7
	75			12	12	11	7	8	5
	50			9	8	7	-	-	2
	25			-	4	-	-	-	-
<i>K. pneumoniae</i>	100	-	2	12	12	12	5	-	7
	75			-	7	9	-	-	2
	50			-	-	4	-	-	-
	25			-	-	-	-	-	-
<i>S. dysenteriae</i>	100	-	1	16	12	8	10	8	4
	75			10	10	6	4	7	4
	50			6	9	6	-	-	-
	25			-	-	-	-	-	-

Keys: LEE (leaf ethanolic extract); SEE (stem ethanolic extract); BEE (bark ethanolic extract); LWE (leaf water extract) SWE (stem water extract); BWE (bark water extract)

Table 3:
Sensitivity of test isolates to some antibiotics

Test isolates	OFL	AUG	TET	AMX	COT	NIT	GEN	NAL	% Resistance
<i>S. faecalis</i>	R	R	R	R	R	R	R	20	87.5
<i>P. mirabilis</i>	11	R	R	R	R	R	R	R	87.5
<i>P. aeruginosa</i>	19	18	20	12	R	12	R	R	37.5
<i>S. aureus</i>	17	R	R	R	R	R	18	R	75
<i>E. coli</i>	R	R	19	R	R	14	16	23	50
<i>K. pneumoniae</i>	R	20	R	R	R	R	17	R	75
<i>S. dysenteriae</i>	R	R	10	R	23	R	R	R	75

KEYS:

OFL- Ofloxacin; AUG- Augmentin; TET-Tetracycline; AMX- Amoxicillin; COT-Cotrimoxazole; NIT-Nitrofurantoin
GEN Gentamycin NAL Nalidixine

Both *F. capensis* plant extracts (100% water and ethanol) had the best inhibition on *P. mirabilis*. This was also much higher than that recorded with the use of established antibiotics of which seven were unable to inhibit the bacterium growth. The zones of inhibition recorded by ofloxacin and gentamycin against *S. aureus* were 17 and 18mm respectively. The extracts also recorded inhibition even at 50% concentration while 6 established antibiotics were unable to inhibit the growth of *E. coli*.

DISCUSSION

These results showed the presence of secondary metabolites in *F. capensis* of which tannins, saponins, alkaloids and phenolics were especially found abundant in all plant parts tested. Even though low at times, the phytochemicals were recorded in all plant parts unlike the findings of Solomon-Wisdom *et al.*, 2011 where certain phytochemicals may be found in only one part of the plant and not in the other. Adeshina *et al.*, 2010, reported that the geographical location of the plant influences the amount of phytochemicals present in it. In this work, the presence of the phytochemicals in all plant parts tested showed that the Oyo State environment favoured the presence of the phytochemicals in the plant. The high phytochemical contents observed in this plant will make it a unique material of medicinal drug screening and research. A high quantity of saponins was recorded especially in its leaf sample, corroborated by Solomon-Wisdom *et al.*, 2011, and contrary to the findings of Oyeleke *et al.*, 2008, which also reported the presence of all the phytochemicals identified except saponins however, the flavonoid content recorded from the leaf sample was slightly lower than that recorded by Adeshina *et al.*, 2010.

Considerable antimicrobial activities of the plant extracts were recorded in this work and these would be as a result of the presence of the bioactive compounds in *F. capensis*. Both the leaf and bark extracts had the highest inhibitory activities against test pathogens while the stem had the least. Ebana *et al.* (1991) reported the inhibition of pathogenic bacteria by alkaloids. The growth of all the pathogens tested for in this work was inhibited by the plant extracts. The high tannin content in the bark would make it a suitable choice especially in treatment of wounds and bleedings (Nguyi, 1988). The extracts had superior microbial inhibitory activities when compared with the antibiotics used and may find application in the management and treatment of the diseases caused by these test

organisms. That the stem ethanol extract recorded microbial inhibition even at 125µg/ml on *E. coli* suggests a strong possibility of the application of very minimal quantities of *F. capensis* as a chemotherapeutic agent in the treatment of skin infections such as pruritis, boils, burns, surgical wound dressings, etc (Akpendu *et al.*, 1994; Solomon-Wisdom *et al.*, 2011).

REFERENCES

- Adeshina G. O., Okeke C. E., Osuagwu N.O., Ehinmidu J. O. (2010): Preliminary in-vitro antibacterial activities of ethanolic extracts of *Ficus sycomorus* Linn. and *Ficus platyphylla* Del. (Moraceae) Afr. J Microbiol. Res. 4 (8), 598-601.
- Ahmadu A. A., Zezi A. U., Yaro A.H. (2007): Anti-diarrheal activity of the leaf extracts of *Daniella oliveri* Hutch and Dalz (Fabaceae) and *Ficus sycomorus* Miq (Moraceae). Afr. J. Trad. CAM. 4 (4), 524-528.
- Amos S., Binda L., Chindo B., Akah P., Abdurahman M., Danmallam H. U., Wambebe C., Gamaniel K. (2001): Evaluation of methanolic extract of *Ficus platyphylla* on gastrointestinal activity. Indian J. Exp. Biol. 39 (1), 63-67.
- Ebana R. U. B., Madunagu B. E., Ekpe E. D., Otung I. N., (1991): Microbiological exploitation of cardiac glycosides and alkaloids from *Garcinia kola*, *Borreria ocymoides*, *Kola nitida* and *Citrus aurantifoli*. J. Appl. Biotechnol. 71, 398-401.
- Hansch C., Taylor J. B., Sammes P. C. (1990): Comprehensive Medicinal Chemistry Vol 1 Pergamon Press. Indiana U.S.A.
- Hassan S. W. (2005): Antimicrobial screening, phytochemical analysis and toxicological studies on some medicinal plants. PhD dissertation, Usmanu Danfodiyo University, Sokoto, Nigeria.
- Irvine F.R. (1961): Woody plants of Ghana with special reference to their uses. PP 347-349. Oxford University Press. London.
- Nguyi A. (1988): Tannins of some Nigerian flora. J. Biotechnol. 6: 221-226.
- Oyeleke, S. B., Dauda, B. E. N., Boye, O. A. (2008): Antibacterial activity of *Ficus capensis*. Afr. J. Biotechnol. 7: 1414 - 1417.
- Sandabe U. K., Kwari H. D. (2000): Some aspects of ethno-Veterinary Medicine among Kanuri and Bura of Borno state Q. J. Borno Museum Soc. pp. 44-45
- Sandabe U. K., Onyelili P.A., Chibuzo G. A. (2006): Phytochemical screening and effects of aqueous extract of *Ficus sycomorus* L. (Moraceae) stem bark on muscular activity in laboratory animals. J. Ethnopharmacol. 104: 283-285.
- Shellard E. J. (1987): Medicines from Plants. Plant Med. 53(2): 6-8.
- Solomon-Wisdom G. O., Shittu G. A., Agboola Y. A. (2011): Antimicrobial And Phytochemical Screening

Activities Of *Ficus Sur* (Forssk). New York Science Journal 4(1):15-18.

Stary F. (1998): Medicinal Herbs and Plants. PP.6-20. Tiger Books International Plc. U.K.

Trease G. E., Evans W. C. (1989): Textbook of Pharmacognosy. 13th Ed. PP. 542-545; W.B. Sanders Company Ltd. London.

Udobi C. E., Onalapo J.A., Agunu A. (2008): Antibacterial activities and bioactive components of the aqueous fraction of the stem bark of *Parkia biglobosa* (JACQ) (Mimosaceae). Nigerian J. Pharm. Sci. 7(1): 49-55.

Wakeel O. K., Aziba P. I., Ashorobi R. B., Umukoro S., Aderibigbe A.O., Awe E. O. (2004): Neuropharmacological activities of *Ficus platyphylla* stem bark in mice. Afr. J. Biomed. Res. 7(2): 75-78.

UNIVERSITY OF IBADAN LIBRARY