
**IBADAN
JOURNAL OF
AGRICULTURAL
RESEARCH**

VOLUME 5 - 9
2009 - 2013

ISSN 0794-5442

UNIVERSITY OF IBADAN LIBRARY

Ibadan Journal of Agricultural Research Volume 5-9

Contents

Page

Vol. 5 – 2009

Interplay of Nutrition and Feeding Frequency on the Growth of African Catfish

Clarias gariepinus Fry

– Akinwale, A. Olusegun, Faturoti, E. Olujimi and Enilolobo, David

3

The Health Situation of Rural Households: A Case of Peri-Urban and Remote Farming Systems in Imo State, Nigeria

– Olajide O. Adeola

8

Vol. 6 – 2010

Benthic Macro-Invertebrates and Physicochemical Quality of Reservoirs in University of Ibadan Fish Farm

– Jenyo-Oni, Adetola, Akinwale, A. Olusegun and Dauda, A. Babatunde

19

Exogenous Oestrogen: Effect of Oestradiol Benzoate on Blood Chemistry of Large White Pigs

– Ladokun A. O. Ladipo M. K. Alabi O. M. Adejumo, D. O.

27

Enhancing Sexual Behaviour in Gilts through the Administration of Testosterone Enanthate

– Ladokun, A.O., Ofite J.R., Sokunbi O.A., Oke O. E. and Adejumo, D O

35

Relative Efficacy of Two Trypanocides on the Treatment of Trypanosome Brucie Infected Grasscutter (*Thryonomys swinderianus*)

– Omonona A.O., Lasisi O.T. and Kolade T.V.

43

Aquatic Tourism Potentials of Okomu National Park, Nigeria

– Ojo, S.O.

49

Vol. 7 – 2011

Agro-Industrial By-products and Farm Wastes for Poultry Production in Nigeria: Opportunities and Challenges

– Abu, O. A. Iyayi, E. A. and Tewe, O. O.

56

Management of Indigenous Pig Breeds in Iseyin, Oyo State of Nigeria

– Adedeji, T.A., Ojedapo, L.O., and Ogundipe, R.I

66

Vol. 8 – 2012

Cytotoxicity Effect and Utilization of *Mytilus edulis* Shell in The Diet of

Clarias gariepinus

– Ayoola, S. O

76

Hepatic Enzyme Profile of Black Jaw Tilapia *Sarotherodon melanotheron*, in Relation to Water Quality in the Lagos Lagoon

– Ayoola, S. O and Bello, M. D

84

Influence of Flake Thickness on Flexural and Moisture Properties of Wood Cement Panels Fabricated from Mixed Furnish of Eight Hardwood Species – Badejo, S. O. and Omole, A. O.	93
Nature Tourism Potential of Bitumen Belt of Ode-Irele, Ondo State, Nigeria – Ayodele, I.A, Ojo, S.O. and Ogunsusi, K.	109
Spatial and Temporal Distribution of Plant-Parasitic nematodes Associated with Banana and Orange from Orchards in Ibadan, Nigeria. – Claudius-Cole, A. O.	121
Vol. 9 –2013	
Cooperative Credit: Effect on Productivity and Welfare of Rural Farming Households in Yewa North Local Government Area of Ogun State, Nigeria – Ambali, O. I. Aminu, F. O. and Ologbon, O. A. C.	130
Meat Quality Characteristics of Broilers Fed Probiotics Feed Supplement – Adebiyi Olufemi A.	142
Soil contamination by lead acid battery wastes inhibits seed germination and induces morphological and physiological changes in seeds of maize (<i>Zea mays</i> L.) and <i>Jatropha curcas</i> . – Adejumo, S. A. and Agunsoye, D. O.	148
Growth and Yield Response of <i>Moringa oleifera</i> (Lam) to Different Rates, Methods and Timing of Compost Application – Adejumo S.A. and Olaleye O. O.	161
Performance and Crude Digestibility in Broilers Fed Diets Formulated Based on Total or Digestible Crude Protein – Agboola, A. F. and Iyayi, E.A.	177
Food and feeding habits of <i>Mugil cephalus</i> (grey mullet) from wetland of Ogun Water-side Local Government Area – Idowu, A.A, Odulate, D.O, Adeosun, F.I, Abdul, W.O, Akinyemi, A.A and Akinware, T.H.	183
Knowledge, Attitude and Practice of Child Nutrition among Women in Southwest Nigeria – Thomas, K.A and Oladeji, J.O	190
Physico-Chemical Properties of Meat and Blood Profile of Broiler Chickens Fed Five Commercial Vitamin-Mineral Premixes in Ibadan, Nigeria – Ogunwole, O. A., Omojola, A. B., Ayinde, B. O., Taiwo, B. A. and Kolade, E. O.	200
Blood Profile of Broiler Chickens as Affected by Diets Supplemented with Graded Levels of Ascorbic Acid – Ogunwole, O. A., Oikeh, I., Majekodunmi, B. C, Ayinde, B. O., and Lawal, T. T.	211

Use of Information Communication Technologies (ICTs) for extension delivery in selected Research Institutes and Agricultural Development Programmes (ADPs) in southwestern Nigeria – Olajide, B. R and Amusat, A. S	218
Sourcing and usage of Standard Cassava Cuttings by Farmers in Egbeda Local Government Area of Oyo State – Olajide B. R and Fawole O. P.	226
Effect of Social Capital on the Technical Efficiency of Arable Crop Farmers in Ibarapa Area of Oyo State, Nigeria – Ologbon, Olugbenga A. Christopher; Idowu S. Dele and Oyebanjo, Olumayowa	234
Climate Change and Its Implications on the Assessment of Planting Opportunities in the Dry Northern Nigeria – Oluwasemire K.O.	244
Allelopathic Effect of Aqueous Shoot and Root Extracts of <i>Alternanthera brasiliana</i> (L.) O. Kuntze on Germination and Seedling Growth of <i>Amaranthus cruentus</i> L. and <i>Zea Mays</i> L. – Oluwakemi Owoseni and Rasheed Olufemi Awodoyin	257
Market Integration of Dry Fish Market in Oyo State Nigeria – Temitayo A. Adepoju and V.O. Okoruwa	265
Price Trend and Analysis of Selected Food Items in Rural and Urban Markets in Oyo State – Anidi Oluwakayode and Olajide O. Adeola	275
Assessment of Growth Performance and Meat Quality of African Giant Snail (<i>Archachatina marginata</i> Swaison) Fed With Mullberry (<i>Morus alba</i>) and Siam-weed (<i>Chromolaena odorata</i>) Leaf Meal Based Diet – Imran, G. T., Bobadoye, A. O., Ogunlase, O. and Bobadoye, B. O.	283
Sensory and Quality Evaluation of Differently Cured Snail Meat – Olusola O.O, Aluko O. T and Oyadeyi, O. S	288

SENSORY AND QUALITY EVALUATION OF DIFFERENTLY CURED SNAIL MEAT

Olusola O.O¹, Aluko O. T¹ and Oyadeyi, O. S²

¹ Meat Science Laboratory, Department of Animal Science, University of Ibadan, Ibadan.

² Department of Animal Health and Production, Oyo State College of Agriculture, Igbo Ora.

Corresponding author: olusolaolubunmi@yahoo.co.uk

ABSTRACT

Sixty four snails of the Species *Archachatina marginata* were shelled and the edible part properly washed with clean water. The meat were randomly allotted to four treatments; Wet curing, Dry curing, Wet curing with injection, No curing (control). The meat assigned to the different treatments were cooked with dry heat (grilling) and kept at ambient temperature of 27°C for 14 days. Physical and chemical changes during curing and processing were observed. The pH, processing yield, salt accumulation and sensory evaluation significantly differed ($P < 0.05$) after two weeks of ageing. Snail meats that were wet cured, dry cured and injected with cure (treatments A, B and C) lost much moisture by dripping and evaporation hence, a significant ($P < 0.05$) weight loss was noted in these treatments. Sensory evaluation done showed that the salt content was moderate across treatments. However, wet cured snail meat was ranked best for flavour, tenderness, juiciness and overall acceptability.

Key words: Snail meat, curing, grilling, pH, processing yield.

INTRODUCTION

Snails belong to the phylum mollusca and are used as food, feed or source of revenue (Owolabi, 2009). Snails constitute the major and cheapest sources of protein in Nigeria (Ademolu *et al*, 2004; Fagbuaro *et al*, 2006). They are eaten by the rich and poor, urban and rural dwellers as well as in all continents of the world (ZASBDC, 2008). Popular species of economic interest are *Helix pomatia*, *Helix aspersa* and the West Africa giant snails: *Achatina fulica*, *Archachatina marginata* and *Achatina achatina* (Cobbinah *et al*, 2008; Owolabi, 2009). Snail meat has a protein content of 18.66 - 20.56% (Okonkwo and Anyaene, 2009) its iron content (45-59mg/kg), low in fat (0.05-0.08%) and contains almost all the amino acids needed for human nutrition (Owolabi, 2009). Snail is high in health benefiting essential fatty acids such as linoleic acids. That is 57% polyunsaturated fatty acids, 15.5% of monounsaturated fatty acids, and 23.25% of saturated fatty acids (Su *et al.*, 2004. In folk medicine, the bluish liquid obtained when the

meat has been removed from the shell is believed to be good for brain growth in infant's development (Owolabi, 2009). It is believed in some quarters that snail meat contains pharmacological properties or value in counteracting high blood pressure, hemorrhoids, constipation, arteriosclerosis, kidney related diseases and suppression of stroke. The high Iron content of snail meat is considered important in the treatment of anaemia and in the past the meat was recommended as a means of combating ulcers and asthma (Abdul Azeez, 2009; Owolabi, 2009).

Curing is the addition of salt and spices to meat, it is an age old act of meat preservation. Besides adding to flavour and taste of the final product, salt is important in the reduction of water activity in meat. The content of salt in sausages, hams, corned beef and similar products is normally 1.5-3%. Solely common salt is used if the cooked products shall have a greyish or greyish-brown colour (Gunter & Peter, 2007). This study was carried out to assess the effect of curing snails using salt in its solid state or brine,

and to check its effect on pH, yield and consumer acceptability of snail meat.

MATERIALS AND METHODS

Source and dressing of snail meat.

Sixty four snails of the species *Archachatina marginata* were purchased from a public market in Ekiti State, Nigeria. The shells were carefully crushed to remove the body from the shell. The edible parts (the foot) was separated from the visceral mass and was washed with potassium alum ($K_2SO_4 \cdot Al_2(SO_4)_3 \cdot 24H_2O$) to remove the slime. Then, they were rinsed with distilled water, mopped free of excess water and weighed. The washed snail meats were randomly allotted across the treatments which were; A: Wet curing, B: Dry curing, C: Wet curing with injection D: No curing (control). Samples were randomly selected from each treatment for organoleptic evaluation, physical and chemical analysis.

Composition of curing brine and mixture: The curing mixture was constituted as follows: 50gm of sodium chloride, 3 gm of sugar and 1gram of additives (0.33gm each of ginger, garlic and white pepper in powdered forms) per liter of water. All the samples (Treatment A-D) were subjected to dry heat treatment using the Crown Star Toaster Grill of the model MC-2012250T. Samples were then aged at room temperature of 27°C for two weeks

Treatments: Each group consisted of sixteen snail meat pieces weighing 830 – 890 grams of fresh snail meat per treatment in a transparent plastic bucket with lid. The treatments were as follows: **Treatment A:** Samples were soaked in curing brine. **Treatment B:** Snail meat samples were dry-salted. The curing mixture was randomly rubbed over each sample. Meat pieces were piled on each other with layers of curing salt between them. **Treatment C:** Meat samples were injected with curing brine at the rate of 8%W/W meat using 5ml syringe with needle. **Treatment D:** The samples were not cured (control), these were refrigerated at 0°C while other samples went through the process of curing. After curing, all treatments were equilibrated for 24 hours at ambient conditions, and were rinsed of excess salt.

Sample preparation: Samples of raw and processed products were separately ground with a meat blender before they were used in analysis.

Evaluation of pH: pH-values were determined by dispersion of one gram sample in 10ml of distilled water using a pH meter (model PHS-25). The pH-values were measured at 25°C.

Measurement of salt Accumulation Values: Salt residue was determined as the chloride using the method described in USDA (2009). 1.5g of ground sample was weighed into 300ml flask with 25ml of 0.1N $AgNO_3$ solution, 15ml of conc. HNO_3 was added and boiled until the meat was digested. $KMnO_4$ was added in small portions while boiling until the solution became colourless. 25ml of H_2O was added to the solution, boiled for 5minutes and cooled to room temperature (27 – 29°C). The flask content was diluted to 150ml with distilled water. 5ml diethyl ether was added and titrated with KSCN solution to a permanent light brown colour. Concentration was calculated as:

$$NaCl \% = \frac{[25.0ml - (ml \text{ KSCN})(R)] (NaNO_3)(5.85)}{\text{sample weight}}$$

$$\text{Where } R = \frac{\text{ml } AgNO_3}{\text{ml KSCN}}$$

Processing yield: Samples were weighed at each processing stage. Data obtained were expressed as a percentage of the initial weight using the formula:

$$\text{Processing yield (\%)} = \frac{\text{Initial weight} - \text{Processed weight}}{\text{Initial weight}} \times 100$$

Hydrated yield: Samples from each treatment was weighed. A volume of distilled water weighing 5 times weight of meat sample was measured and boiled at 100°C for 10 minutes. The source of heat was removed and the meat was left in the hot water for 10 minutes. Then the sample was removed from the hot water, moped dry and weighed.

$$R_{ii} \% = \frac{Hw - Dw}{Dw} \times 100$$

Where: Hw= hydrated weight; Dw= Dehydrated weight of dried meat; R_{ii} = Rehydrated percentage

Organoleptic Evaluation: Sensory evaluation was carried out after two weeks aging using a 9 point hedonic scale with 9 as maximum score and 1 the least score. The selected samples were hydrated, cut into bits and served to the panelists for evaluation

Statistical analysis: Analysis of variance was carried out using Completely Randomized Design according to the methods of SAS (1998).

RESULTS AND DISCUSSION

Results obtained are presented in tables 1 to 3. The pH of the fresh snail meat was 7.40- 7.41 for all treatments at onset, ranging near neutrality.

Curing and pH affect in particular the water holding capacity of meat samples. Significant differences were observed in pH values obtained across the treatments at different stages of processing viz curing, grilling and ageing. The trend was similar to that observed by Thorarinsdottir *et al.*, (2001) who reported the pH in fish muscle to have decreased during salting, a rapid drop was observed after dry salting, but a rise in pH during ageing. These changes in pH while processing might be due to protein conformational changes. The high moisture reduction in 'A' might be due to its low pH value. The meat with high pH had better water retention properties (FAO, 1990).

Table 1: Salt Content and pH of Snail Meat

Parameters	A	B	C	D	SEM
Salt content (%)	8.06 ^a	7.79 ^{ab}	7.17 ^b	7.17 ^b	0.14
pH :					
Fresh snail	7.41	7.40	7.41	7.40	0.00
Cured meat	6.62 ^d	7.92 ^b	7.26 ^c	8.11 ^a	0.13
Grilled meat	5.70 ^c	7.25 ^a	8.01 ^b	8.01 ^b	0.14
Aged meat	7.40 ^d	8.03 ^c	8.13 ^b	8.14 ^a	0.07

abc: Mean on the same row with different superscript are significant ($P < 0.05$)

Snail meat naturally contains sodium ions (7.17%) as indicated in table 1. As reported by Maha and James (2003), excess salt has undesirable effects on the flavor, colour and appearance of meat. Application of dry salt can result in unattractive and darker lean. The increased concerns regarding high salt intake and its relative role to hypertension, high blood pressure, and potential heart disease, public health authorities recommend reduction in salt intake. This makes snail meat which has 15% protein, 80% moisture, 2.4% fat (75% of which is unsaturated) an ideal meat for all ages. The weight loss in the meat samples were significantly different ($p < 0.05$). Samples treated with the curing salt had lower moisture content

hence the weight loss of 33.85%, 29.89% and 27.67% respectively after the curing process. As reported by Okonkwo and Anyaene (2009), the equilibration in salt caused the reduction in weight probably due to osmotic dehydration. Though there was significant loss of weight across the treatment on the application of heat by grilling, the result showed treatments D and B lost more weight (54.84% and 41.58% respectively). The reduced moisture suggests decrease in water activity and increasing microbiological safety and stability especially through inhibition of bacterial organisms (Thorarinsdottir, 2001). Also, there was no significant difference in rehydration property. This implies the meat samples were evenly dried.

Table 2: Processing yield

Parameters	Wet cure	Dry cure	Wet-cure injected	control	SEM
Weight loss (%):					
In cured snail	27.67 ^a	33.85 ^a	29.89 ^a	14.84 ^b	2.32
In grilled snail	41.58 ^{ab}	30.27 ^b	37.32 ^b	54.84 ^a	3.29
In aged snail meat	52.24 ^a	34.00 ^b	31.11 ^b	37.42 ^{ab}	3.24
Rehydrated %	31.63	27.52	31.47	35.99	

abc: Means on the same row with different superscript are significant ($P < 0.05$).

Table 3: Sensory evaluation rating of cured meat aged for 14 days

Parameters	Wet cure	Dry cure	Wet-cure injected	Control	SEM
Colour	2.33	2.25	1.87	1.87	0.09
Flavour	4.33 ^a	4.12 ^a	3.22 ^b	3.31 ^b	0.16
Tenderness	4.04 ^a	3.50 ^a	3.58 ^a	2.39 ^b	0.21
Juiciness	3.46 ^a	3.29 ^{ab}	2.79 ^{bc}	2.25 ^c	0.16
Saltiness	4.22 ^b	3.00 ^c	6.69 ^a	3.00 ^c	0.46
Overall	5.12 ^a	5.12 ^a	4.25 ^c	4.59 ^b	0.12
Acceptability					

abc: Means on the same row with different superscript are significant ($P < 0.05$).

Fresh snail meat is dark, salting and ageing makes it even darker. All samples were scored as extremely dark. Wet cured meat ranked highest for flavour, tenderness, juiciness and overall acceptability. Wet cured meat with injection was saltiest probably due to brine injection,

CONCLUSION

With the fast growing population of the urban areas, the demand for meat is on the increase. Heliculture is one of the means to meet the increasing protein demand. Wet curing of snail meat in salt and spices for 24 hours before grilling will enhance its organoleptic traits as well as its keeping quality at ambient temperature.

REFERENCES

- AbdulAzeez, I. (2009). Medicinal value of snails in Nigeria. Article base. www.articlebase.com
- Ademolu K.O. Idowu A.B., Mafiana C.F., Osinowo O.A (2004). Performance, proximate and mineral analyses of African giant land snail (*Archachatina marginata*) fed different nitrogen sources. *African J Biotech* 2004; 3(8): 412-417.
- Cobbinah, J.R., Vink, A., Onwuka, B. 2008. Snail farming; Production, processing and marketing. Agromisa Foundation, Wageningen
- F.A.O (1990): Manual of Simple Methods of Meat Preservation. FAO Animal production and health paper no. 79, Rome, FAO.
- Fagbuaro, O., Oso, J.A., Edward, J.B., and Ogunleye R.F. (2006). Nutritional status of four species of giant land snails in Nigeria. *Journal of Zhejiang University Science* 7(9):686-689
- Gunter, H., Peter, Hautzinger. (2007). Meat processing technology for small-to medium-scale producers. FAO- the United Nations regional office for Asia and the Pacific, Bangkok. Pp 59-171.
- Institute Food Technologist. (1980). Dietary salt: a scientific status summary. *Food technology* 33:85-91.
- Maha N. H and James L. M (2003). Salted meat. *Food and culture Encyclopedia Microsoft Encarta 2009. 1993-2008 Microsoft Corporation.*

- Okonkwo, T.M. and Anyaene, L.U. (2009). Meat yield and the effects of curing on the characteristics of snail meat. *Journal of Tropical Agriculture, Food, Environment and Extension*. 8(1):66-73.
- Owolabi, M.F. (2009). Snail farming and management. www.efarmspro.com
- SAS (1998). Statistical Analysis System Institute's User's guide SAS Institute Inc. NC.54:365-367
- Su, X.Q., Antonas, K.N., and Li, D. (2004). Composition of n-3 polyunsaturated fatty acid contents of wild and cultured Australian abalone. *International Journal of Food Sciences and Nutrition*, 55(2): 149-154.
- Thorarinsdottir K.A., Arason S., Bogason S.G., and Kristber K. (2001). Effects of Phosphate on Yield, Quality and Water-holding capacity in the Processing of salted Cod (*Gadus morhua*) *Journal of Food Chemistry and Toxicology* Vol. 66, No.6.
- United States Department of Agriculture Food Safety and Inspection Service, office of Public health Science. 2009. Determination of salt.
- ZASBDC. 2008. Canned snail meat- China canned food, seafood, meat in canned food. Zhejiang Aotokang Special Biology Development Centre [ZASBDC], China

Parameter	Value	Value	Value
Color	4.33	4.33	4.33
Flavor	4.33	4.33	4.33
Tenderness	4.33	4.33	4.33
Moisture	4.33	4.33	4.33
Water-holding capacity	4.33	4.33	4.33
Acceptability	4.33	4.33	4.33

Table 1: Means and standard deviations for different attributes of snail meat.

Fresh snail meat is dark, salty and aging makes it even darker. All samples were scored as extremely dark. Wet cured meat ranked highest for tenderness, firmness and overall acceptability. Wet cured meat with injection was highest probably due to its injection. The injection probably due to its injection.

CONCLUSION
With the fast growing population of the urban area, the demand for snail as an alternative protein source is increasing. Wet curing of snail meat will be an effective way to meet the increasing protein demand. Wet curing of snail meat will be an effective way to meet the increasing protein demand. Wet curing of snail meat will be an effective way to meet the increasing protein demand.

REFERENCES
Abdulkareem, I. (2009). Medicinal value of snail in Nigeria. *African basic*. www.ajbas.ac.uk

Adeniyi, K.O. Ibadan A.B. Martins C.F. (2004). Performance, proximate and mineral analysis of African giant land snail (*Archaiatus marginatus*) fed different nitrogen sources. *African Journal of Biotechnology* 3:413-417.

Coplin, R. Vink A. Olowu E. 2008. Snail meat production, processing and marketing. *Agriculture Foundation*, Washington, D.C.

F.A.O. (1990). Manual of snail production and health. *FAO*, Rome.

Richard O. Ode. 2008. Snail meat and its nutritive value. *Journal of Animal Production* 4:1-4.

Ogunleye, R.E. (2006). Nutritional value of four species of giant land snails in Nigeria. *Journal of Nigerian University Science* 1(1):1-10.

Quar, H. Peter. 2008. Snail meat processing technology for small-scale producers. *FAO*, Rome.

National compound order for Asia and the Pacific. *Bangkok* 171.

Institute Food Technology (1980). *Dietary snail: a scientific status summary*. Food technology 31:52-61.

Mahn N. H and James E. W. (2003). Salted meat. *Food and animal biotechnology* 19:1-10.

Environ 2009. 1993. *Environ* Corporation