

## Bilateral congenital cystic adenomatoid malformation of the lungs : a case report

OM Atalabi\*, AO Ogunseyinde\*, MO Obajimi\*, AG Falade+, AO Adebosè, and IE Jite+

Departments of Radiology\*, Paediatrics+ and Surgery±, College of Medicine, University College Hospital, Ibadan, Nigeria

### Summary

A six weeks old infant presented with scalp ulcer and fever, and on examination was found to have resonant percussion notes bilaterally. The initial chest radiograph revealed multiple lucencies which were initially thought to be due to diaphragmatic hernia, but the dilemma was resolved by Computerised tomography which revealed the lucencies to be multiple cysts characteristics of Congenital Cystic Adenomatoid Malformation (CCAM) type II.

**Keywords:** Congenital cystic adenomatoid malformation, radiological diagnoses.

### Résumé

Un bébé de six semaines avait l'ulcère du crâne et la fièvre. A l'examen clinique, il a été observé qu'il avait des notes de résonance de percussion bilatérale. Un examen de la radio révélait des multiples traces qui étaient initialement pensées dues à l'hernie diaphragmatique. Le dilemme était résolu par la tomographie informatisée qui révélait que ces traces étaient des kystes caractéristiques d'une malformation kystique adénomatique congénitale de type II.

### Introduction

Congenital cystic adenomatoid malformation (CCAM) is a developmental hamartomatous abnormality of the lung with adenomatoid proliferation of cysts resembling bronchioles. Ch'in and Tang first described this disease entity in 1949. CCAM represents approximately 25% of all congenital lung lesions [1]. Although the cause is not known, CCAM is believed to result from focal arrest in fetal lung development before the seventh week of gestation secondary to a variety of pulmonary insults. Most infants born with CCAM have no symptoms at birth. Many women in the developing world do not have access to routine prenatal ultrasound scan, therefore CCAM is rarely diagnosed de novo except accidentally as in this case report; as an extensive literature search had not revealed any reported case of CCAM in this environment.

### Case report

A.O was a female infant admitted at six weeks of birth with scalp ulcer and fever of twelve and ten days duration respectively. The patient was well until about two weeks

before presentation when the mother noticed that the baby was having low-grade fever and two days later noticed an area of scalp discoloration over the occipital region, which was discharging purulent materials with associated sloughing of the overlying skin. She had received treatment at home with antibiotics and other unknown drugs given by the mother. The baby was born at term and the pre- and post-natal history was essentially normal with no ingestion of any unusual drug during pregnancy by the mother. On physical examination, patient was found to be acutely ill, pale, anicteric, acyanosed and dehydrated. In addition she had bilateral pitting pedal oedema up to the ankle, worse on the left and a tender swelling on the right middle finger. She had a scalp ulcer, extending from the right parieto-temporal region across the midline involving the whole occipital region, with surrounding necrotic skin. The floor of the ulcer was hard, with an island of bone at the centre measuring approximately 6cm x 5cm.



Fig. 1:

Chest percussion was resonant bilaterally with vesicular breath sounds; the pulse was irregular at 164/minute. There was no cardiac murmur. The antero-posterior chest radiograph showed bilateral areas of large cystic lesions, devoid of lung markings with surrounding septated walls one of which showed a fluid level on the left upper zone. The lateral film showed these large cystic lesions to be mostly posterior, suggesting a diaphragmatic hernia (Fig. 1). However a limited barium meal



study revealed no abnormality of the gastrointestinal system. Computerised tomography of the chest showed multiple well-defined cystic lesions of varying sizes in both lung fields, with the largest measuring 4.2 x 3.5 cm (Fig. 2). Based on the overall radiographic appearances a diagnosis of congenital cystic adenomatoid malformation type II was made.

The patient however died before any surgical resection could be performed.

anomalies, and the prognosis is poor [2]. The type III lesion is a large, bulky non-cystic lesion producing mediastinal shift. Bronchiole-like structures are lined by ciliated cuboidal epithelium and separated by masses of alveolus-sized structures lined by non-ciliated cuboidal epithelium [3, 4].

Majority of patients present as respiratory emergencies in the newborn although the disease may remain undetected for some years. Diagnosis is usu-




Fig. 2:

### Discussion

Congenital cystic adenomatoid malformation (CCAM) is a developmental hamartomatous abnormality of the lung with adenomatoid proliferation of cysts resembling bronchioles, and accounts for approximately 25% of all congenital lungs [11]. Stocker classified congenital cystic adenomatoid malformation (CCAM) into three types based on clinical, gross and microscopic criteria. Type I lesion is composed of single or multiple large cysts (more than 2 cm. in diameter), frequently producing mediastinal herniation. The cysts are lined by ciliated pseudostratified columnar epithelium. The walls of the cysts contain prominent smooth muscle and elastic tissue. Mucus producing cells are present in approximately one-third of the cases, and cartilage in the wall is rarely seen. Relatively normal alveoli may be seen between the cysts. The type II lesion is composed of multiple small cysts (less than 1 cm. in diameter) lined by ciliated cuboidal to columnar epithelium. Structures resembling respiratory bronchioles and distended alveoli are present between the epithelium-lined cysts. Mucous cells and cartilage are not present. Striated muscle fibres may be seen rarely. The type II lesion is associated with a high frequency of other congenital

ally made prenatally using ultrasound, and postnatally using chest radiograph and computerised tomography (CT). On chest radiograph, the malformation is usually composed of air filled cysts of varying sizes which may contain fluid levels, while on CT, air filled cysts of varying sizes which may contain fluid are seen [5- 8], and these were demonstrated in our patient Figs 1 and 2. In the study done by Hubbard *et al.*, [9] MRI is useful as adjunct to prenatal diagnosis of CCAM thus allowing for choice of treatment and parental counselling. Khosa J.K. *et al* in their study also concluded that CT scanning is mandatory for postnatal evaluation because chest radiograph could be normal [10]. The major diagnostic differential is diaphragmatic hernia, which can produce a similar multicystic appearance on plain chest x-ray, however in CCAM there is normal bowel gas pattern [5].

Due to an increasing trend in the detection of asymptomatic antenatally diagnosed CCAM, consideration is given to early surgical excision to prevent complications and recurrence [9]. Prognosis is based on findings of prenatal ultrasound and it has been found that microcystic lesion, bilateral lung involvement and hydrops were each highly correlated with poor prognosis.

while neither polyhydramnios nor mediastinal shift was significantly associated with bad outcome [4].

#### References:

1. Gerald Mandell, congenital adenomatoid cystic malformation: e-Medicine articles.
2. Bunduki V, Ruano R, da Silva MM, *et al*: Prognostic factors associated with congenital cystic adenomatoid malformation of the lung. *Prenat. Diagn.* 2000; 20: 459-464
3. Stocker JT, Madewell JE and Drake RM: Congenital cystic adenomatoid malformation of the lung. Classification and morphologic spectrum. *Hum Pathol.* 1977; 155-171
4. Kim WS, Lee KS, Kim IO, Suh YI, *et al*: Congenital cystic adenomatoid malformation of the lung: CT pathologic correlation: *Am J Roentgenol.* 1997; 168: 47-53
5. Grainger RG and Allison DJ. *Diagnostic Radiology* 1992. Second Edition, Volume 2, Churchill, Livingstone, London. 380.
6. Laberge JM, Flageole H and Pugash D. Outcome of the prenatally diagnosed congenital cystic adenomatoid lung malformation: a Canadian experience. *Fetal Diagn Ther.* 2001; 16: 178-186
7. Lane F.D. *Fundamentals of Paediatric Radiology*. First edition, WB Sanders Company pg.35.
8. Marshall KW, Blane CE, Teitelbaum DH, *et al*: Congenital cystic adenomatoid malformation: impact of prenatal diagnosis and changing strategies in the treatment of the asymptomatic patient. *AJR Am J Roentgenol* 2000; 175: 1551-1554
9. Hubbard AM, Adzick NS, Crombleholme TM, *et al*: Congenital chest lesions: diagnosis and characterization with prenatal MR imaging. *Radiology* 1999; 212: 43-48;
10. Khosa J K, Leong SL, Borzi PA. Congenital Cystic Adenomatoid Malformation of the lung: *Pediatr Surg Int.* 2004; 20: 505-508

Received: 11/07/05

Accepted: 04/07/06

UNIVERSITY OF LEADAM



## Notes for Contributors

All manuscripts should be addressed to the Editorial Office of the African Journal of Medicine and Medical Sciences, Institute for Advanced Medical Research and Training (IMRAT), College of Medicine, University College Hospital, Ibadan, Nigeria.

Manuscripts are accepted subject to the understanding that no substantial part has been or will be published elsewhere. This does not refer to abstracts of oral communications that are printed in proceedings of societies or symposia.

Authors should send three complete copies of the manuscripts (maximum of 18 quarto-sized typed pages including bibliography, illustrations and tables) and retain one copy for reference. Apart from the three typescript copies on white bond paper, submission of a 3.5" diskette is required at acceptance stage. Authors should ensure that the diskette is clearly labelled with the paper title and name(s) of author(s). The diskette must be IBM compatible on Word Processing format up to version 6.1.

Manuscripts being submitted must be accompanied by a covering letter affirming that the paper is submitted only to this Journal. In the case of a paper with more than one author, all must sign the covering letter confirming that they participated sufficiently in the work.

A non-refundable processing fee of N1000.00 (£50.00, \$80.00) per article will be charged authors who do not subscribe to the Journal while subscribers will pay N500.00 only. In addition, authors of papers submitted from Nigeria should send 4 large and 2 officials size envelopes with N400.00 worth of postage stamp for subsequent correspondence. The Editor does not accept responsibility for damage or loss of papers submitted.

Manuscripts, in English Language, should be submitted typed double-spaced and with at least 2.5 cm margins around.

On a single separate sheet, there must be the following: (a) title, (b) author's name and initials, (c) department/s in which the work was done, (d) the name and address of the author to whom correspondence should be addressed, (e) author's present address if different from the department/s in which the work was done, (f) if paper was presented at a meeting, please indicate name of organization, city month and year.

Titles should be short, specific and clear. Omit phrase such as "The use of, "observations on". Authors should provide a shorter running title and up to six keywords. Manuscripts must be accompanied by a summary not exceeding 200 words. The summary will be translated into French by the journal office at a cost to the author. Currently, the fee for translation is N750.00 (\$30.00) non-refundable, which shall be forwarded by a bank draft to accompany the manuscript when found acceptable for publication. Figures and photographs must be submitted as glossy prints, untrimmed and unmounted. Each should be numbered with pencil lightly on the back with the top indicated. The numbers of photographs and illustrations should be kept to a minimum. The legends for figures and tables that should be numbered with Arabic numerals should appear on a separate page.

The author must pay for publication of colour transparencies submitted at the time of acceptance.

All details on charts and graphs must be clearly legible when reduced to the size used in this journal.

The onus of preparing a paper in a suitable form for publication rests with the author. The need for editorial revision for badly prepared typescripts or diagrams may lead either to rejection of the article or delay in publication.

Authors should indicate by a statement in the body of their paper that they complied with the standard requirements of the Ethics Committee of the institution in which the work was done. A Letter of Ethics Committee approval must also accompany the manuscripts at the time of submission. Where an Ethics Committee is not readily available, the Helsinki Declaration principles as revised should be followed strictly.

Workshop and conference reports should not exceed 3 to 10 double-spaced quarto-sized pages. Viewpoints which could be papers expressing personal or group opinion on political, socio-economic and other matters as they relate to the practice of medicine should be limited to 10 A4-sized typed pages. Letters to the Editor may be comments on papers published in the Journal or clinical observations, replies to comments, or other matters of importance and relevance to medicine and related professions. It should not exceed 500 words with a few references and one or two tables and figures.

This Journal has agreed to accept manuscripts prepared in accordance with the Vancouver style and the Editor will consider only papers conforming to this style.

References should be numbered in the order in which they are cited in the text. At the end of the article, the references should be listed as numbered in the text. Each reference should give the names and initials of all authors (unless there are more than six, when only the first three should be given, followed by *et al.*). The authors' names should be followed by the title of the article, the journal title (abbreviated according to the style of *Index Medicus*), the year of publication, the volume number and the first and last page numbers. Titles of books should be followed by the publisher, place of publication, and year. Examples of format for references are as follows:

Edington JM, Osunkoya BO, Smith JA.

Immunopathology of Burkitt's lymphoma. West Afr Med J 1986; 85: 76 - 87.

Brown A. Primary Health Care and the Medical Curriculum. Edinburgh: Universities Press. 1977.

Lewis A. Primary liver cell carcinoma. In: Ajose A, Odeku EL, Eds. Priorities in Health Planning. Ibadan: University Press, 1983; 110 - 117

Reference to tables should be in Arabic numerals, e.g. Table 3, and tables should include titles, which make them meaningful clear without reference to text. Tables should be typed separately from the text.

Proof corrections are expensive and correction of proofs other than printers' errors should be kept to a minimum. Authors must return proof corrections within 3 days of receipt. Failure to do this will result in the paper being published with the Editor's corrections only. Papers accepted for publication remain the copyright of the Journal.

Offprint may be ordered directly from the Editorial office of the Journal. This must be paid for at the time of final acceptance of paper.

Four journal pages will be printed at no cost to the author. Extra pages will attract page-printing charges at the rate of N500.00 (£5, \$7) per page. (approximately 5 quarto-size pages of manuscript).