



ISSN: 0303-3872

**AFRICAN**  
**JOURNAL OF EDUCATIONAL**  
**RESEARCH**

VOLUME 26 JUNE, 2022

JOINTLY PUBLISHED BY DEPARTMENTS OF:  
ARTS AND SOCIAL SCIENCES  
EDUCATION,  
EARLY CHILDHOOD AND  
EDUCATIONAL FOUNDATIONS AND  
SCIENCE AND TECHNOLOGY EDUCATION,  
UNIVERSITY OF IBADAN.

**AFRICAN JOURNAL OF EDUCATIONAL  
RESEARCH**

**Vol. 26, JUNE, 2022**

**Jointly Published By Departments of:  
Arts and Social Sciences Education,  
Early Childhood and Educational Foundations, and  
Science and Technology Education,  
University of Ibadan.**

UNIVERSITY OF IBADAN LIBRARY



## TABLE OF CONTENTS

<b>Student Satisfaction Experience and Infrastructural Facilities on Computer Based Test, in Tertiary Institutions in Lagos State, Nigeria</b> <i>Adebowale Adebagbo, Ph.D and Lucas B. Ojo, Ph.D</i>	1 - 6
<b>Age, Television Viewing and Students' Academic Achievement in Sagamu, Ogun State, Nigeria: Implications for Social Welfare Counselling Practice</b> <i>T. A. Adebowale, Ph.D and Orekelewa Olukayode</i>	7 - 12
<b>Physics Teachers' Perception and Utilization of ICT Tools for Teaching Senior School Physics in Sokoto, Nigeria</b> <i>Afees Akanni Amuda and Ridwan Enuwa Mohammed Ph.D</i>	13-18
<b>Drug Compliance and Psychosocial Wellbeing of Patients with Mental Illness in Ibadan</b> <i>Abimbola Afolabi Ph.D</i>	19-24
<b>English Language Teachers' Job Satisfaction and Commitment to Teaching in Selected Secondary Schools in Ado Ekiti, Nigeria</b> <i>Christianah Adejokemi Daramola</i>	25-33
<b>Awareness and Utilization of Technology-Based Instructional Resources (Tbir) among Business Educators in Colleges of Education in Edo and Delta States of Nigeria</b> <i>Liadi, Hakeem O. Ph.D., Adedapo, A. Ph.D. and Prof. Ibrahim O. Salawu</i>	34-41
<b>Self-Efficacy, Parental Involvement and School Adjustment of High Achieving Secondary School Students in Oyo State, Nigeria</b> <i>Adelodun G. A. Ph.D and Salako, Adebayo Adeyinka Ph.D</i>	42-47
<b>Cultural Content of Intensive English and Effective English Textbooks</b> <i>Gideon Mojere</i>	48-55
<b>Direct and Indirect Comprehensive Metacognitive Feedback Strategies and Senior Secondary School Students' Achievement in English Composition in Ondo State, Nigeria</b> <i>Kehinde Olufemi Ogunyemi and Oluwasuyi Samuel Yakubu</i>	56-62
<b>COVID-19 Pandemic: Response and Recovery among Adult English Language Students in Southwest Nigeria</b> <i>Oladipo Adeyeye Olubodun, Ph.D</i>	63-68
<b>Family Resources, Peer-Rejection, Academic Stress and Locus of Control as Predictors of Drop-out Risk among In-School Adolescents in Ibadan</b> <i>Abulfatai Adekunle Owodunni</i>	69-75
<b>Ibadan Residents' Urban Renewal Knowledge and Attitude: A Qualitative Study</b> <i>Moses Aderibigbe Oyekanmi, Ph.D</i>	76-83
<b>Development and Validation of Students' Evaluation Scale for Lecturer's Teaching Effectiveness in Colleges of Education.</b> <i>Ojo, Oloyede Akinniyi</i>	84-91
<b>Refocusing Teaching and Learning through Reflection for Students' Conceptual Change in Biology in Secondary Schools in Abeokuta, Nigeria</b> <i>Prof. Ige, Temisan A., Prof. Olagunju, Alice M., Oke, Olubukola Abidemi Ph.D and Erinfolami, Oluwatosin Bimbo Ph.D</i>	92-97
<b>Exploring the Impact of Entrepreneurship Education on Enterprise Creation of Undergraduate Students in South-West of Nigeria</b> <i>Ayo-Sobowale, Mopelola Omotayo Ph.D</i>	98-102
<b>Customer Profiling and Utilization of Electronic Banking Services among Motor Parts Traders in Akwa Ibom State</b> <i>Comfort Idongesit Charles and Otobong Udo Etuk</i>	103-107
<b>Influence of Mass Media on Knowledge and Awareness of Covid-19 Pandemic and its Preventive Measures among Residents in Lagos Metropolis</b> <i>Aaron Akinloye</i>	108-112
<b>Child-Friendly School as a Predictor of Pupils Socio-Personal Lives in Oyo Metropolis, Nigeria</b> <i>Oluwatoyin Abike Odeleye, Ph.D, Leye-Akinlabi, Gbemisola and Gbadeyanka, Taofeek Akolade</i>	113-116
<b>Relationship between School Facilities and Academic Performance of Senior Secondary School Students in Ifo Local Government Area, Ogun State, Nigeria</b> <i>Oba-Adenuga, Olusegun Adeleke, Ph.D</i>	117-122
<b>Ekiti State University Science Education Lecturers' and Science Education Undergraduates' Attitude Towards Teaching Practice</b> <i>Raimi, S. M.</i>	123-128
<b>Functional Business Education Programme: A Sustainable Approach to Economic Security in Nigeria</b> <i>Seyi David, Ph.D and Deborah Olufunke Ayeni, Ph.D</i>	129-133
<b>Disease Education on Knowledge, Attitude and Prevention of Measles among Mothers of Under-Five in Ibadan Metropolis, Nigeria</b> <i>S. A. Famuyiwa, Ph.D, Adebayo David Oluwole, Ph.D and L. O. Lawa!</i>	134-142
<b>Social Support and Quality of Life among the Elderly in Ibadan Metropolis</b> <i>Fatoye Helen Ajibike</i>	143-149



# Drug Compliance and Psychosocial Wellbeing of Patients with Mental Illness in Ibadan

Abimbola Afolabi Ph.D

Department of Social Work, Faculty of Education, University of Ibadan, Nigeria.

## Abstract

The study looked at medicine adherence and psychosocial well-being in patients with mental illnesses at Adeoyo State Hospital, Ring Road, Ibadan. A descriptive survey design was used in the investigation. One hundred and twenty (120) people with mental illnesses who visit Adeoyo State Hospital, Ring Road, Ibadan, were purposefully chosen. For data gathering, a customised questionnaire was employed. Data for demographic parameters were analysed using percentage distribution and frequency counts, and Pearson Product Moment Correlational Analysis (PPMC) was used to test hypotheses at  $p = 0.05$ . The study found a strong link between medication compliance and psychological well-being in mentally ill individuals ( $r = .275$ ,  $n = 120$ ,  $p (.002).05$ ). It was also shown that there was a substantial association between medication compliance and respondents' social well-being ( $r = .327$ ,  $n = 120$ ,  $p (.001).05$ ). It was recommended that patients with mental illnesses be monitored to ensure drug compliance. Medical social workers are needed to educate the general population and society about the consequences of drug noncompliance.

**Keywords:** Patients, Drug compliance, Mental illness, and Psycho-social wellbeing.

## Introduction

There is mounting evidence of mental illnesses worldwide. Mental health issues are one of the most significant contributors to the global disease and disability load. Mental health issues account for five of the top ten primary causes of disability worldwide. They are as significant in low-income nations as they are in wealthy ones, spanning age, gender, and socioeconomic class. Furthermore, all estimates imply that mental health problems will skyrocket in the future (Brundtland 2000). Mental illnesses are conditions that have an impact on a person's mood, thoughts, or behaviour. Serious mental disorders include schizophrenia, bipolar disorder, panic disorder, obsessive-compulsive disorder, and major depressive disorder, among others. Although these conditions might be frightening, it is vital to realise that they are curable. Individuals with various disorders can live full, fulfilling lives if they get treatment as needed. Being diagnosed with a major mental illness might come as a surprise to both the individual being diagnosed and his or her family and friends. On the other hand, obtaining a diagnosis and treatment plan may sometimes assist in easing family tension and beginning the healing process.

Noncompliance with antipsychotic medication has a deleterious influence on the course of disease, increasing the likelihood of relapse, re-hospitalization, and suicide, as well as increasing healthcare expenses. Noncompliance with psychiatric drugs or treatment regimens has a significant impact on the progression of mental illness, relapse, and future recovery. Furthermore, there may be a major influence on the expense of treatment, as well as significant barriers to the patient's long-term adaption, such as social, occupational, and academic functioning. Patients suffering from psychological ailments have higher levels of noncompliance than those suffering from physical illnesses. Biological measurements, clinician ratings, patient self-report, pill count, caregiver reports, and medication side effects can all be used to assess treatment adherence. Issues connected to the therapy, patient-related factors, health care, and

socioeconomic situations have all been listed as reasons for non-adherence.

Mental health is an important component of overall health and well-being, but it has received little attention in this region of the world. Furthermore, the World Health Organization (WHO) estimates that 450 million people worldwide have a mental problem, and that 25% of the population will experience mental disease at some point in their life (WHO, 2019). According to the World Health Organization, mental health is "a condition of well-being in which the individual recognises his or her own potential, can cope with the usual demands of life, can work successfully and fruitfully, and can contribute to his or her community" (WHO, 2018).

Various words have been used to characterise the behaviour of continuing to take prescription medicine. The most often used, conventional phrase is "compliance," which has been described as the degree to which a consumer's behaviour corresponds to the prescriber's instructions (Horne, Weinman, Barber, Elliot, & Morgan, 2020). The term "compliance" has been described as "the extent to which a person's behaviour corresponds with medical or health recommendations" (Haynes, 2021) or "the degree of conformity between treatment behaviour and treatment standards" (Hunt, Bergen, and Bashir, 2002). The first of these meanings implies paternalism on the side of the doctor, but the latter, more contemporary term, acknowledges the patient's responsibility in treatment decisions (Hunt, Bergen and Bashir, 2020). The phrase "compliance" is becoming less popular since it indicates a lack of customer engagement and, instead implies a passive strategy in which the consumer dutifully (and often unquestioningly) follows the advice and directives of the healthcare practitioner (Horne et al., 2005; Swaminath, 2017). The premise that medical counsel is helpful for the customer and that rational consumer behaviour entails precisely following medical advice is inherent in many definitions of compliance (Swaminath, 2017).

Schizophrenia is a serious mental condition that often has substantial consequences for people's



quality of life. The amount to which people with schizophrenia take their antipsychotic drugs is thought to have a significant impact on their results. While medication adherence in people with schizophrenia has been widely examined, the majority of research has been quantitative, and therefore the opinions of consumers have been largely ignored. One reason for this omission has been proposed: the belief that people with schizophrenia would be unable to contribute meaningfully to knowledge. It would be preferable to gather more information from patients in order to understand why they do or do not take their drugs.

When compared to non-compliance, medication compliance in patients with schizophrenia is associated with improved results in terms of symptom intensity, relapse, and overall functioning. Several studies have found that non-compliant people with schizophrenia have more severe disease symptoms. Extreme symptom exacerbations frequently result in a recurrence of psychosis and hospitalisation for noncompliant clients. A three-year study of outpatients with schizophrenia indicated that symptom remission was more common among consumers who were compliant with their medication at the time of follow-up (Novick et al., 2010). On the other hand, Rosa, Marcolin, and Elkis (2005), on the other hand, discovered that non-compliant customers experienced an initial exacerbation of symptoms that remained stable during a one-year follow-up. Furthermore, Janssen et al. (2006) discovered that the absolute difference in Positive and Negative Syndrome Scale (PANSS) scores measured at admission and at discharge was significantly greater for patients who were more compliant in their study comparing symptom severity among hospitalised consumers. Medication adherence has been demonstrated repeatedly to minimise the high rates of relapse in patients with schizophrenia (Novick et al., 2010).

Medical intervention in the form of antipsychotic medication is now widely accepted as the primary therapy for schizophrenia (Noetzel, Jones, and Conn, 2012). While antipsychotic drugs are extremely beneficial to the majority of people with schizophrenia, they are not a cure. In many cases, antipsychotic drugs ease or milder symptoms and, in certain situations, can reduce the duration of a schizophrenic episode. The most frequent method of administering antipsychotic medicine is in tablet or liquid form, one to three times per day. Some antipsychotic drugs are also available in tablet form, as well as dissolvable forms (Mueser & Gingerich, 2006). Antipsychotic drugs can also be administered as short-acting and long-acting injections.

Myers and Branthwaite randomly assigned patients in an early trial at a psychiatric outpatient clinic in the United Kingdom into groups that got therapy once a day, three times daily, or selected one of the two schedules. A pill count and an interview were used to determine compliance. There was no overall difference in reported compliance between patients receiving once-day or three-times-daily dosages, although those

who chose to take their medicine three times daily reported higher compliance.

Lin found that patients were more likely to continue taking their medication during the first month of treatment if they had received specific educational messages, such as the fact that they should take their medication daily, that they may not notice any benefit for the first 2–4 weeks, that they should continue even if they felt better, and that they should not stop medication without consulting their doctor. They were also given advice on how to get answers to medication-related questions.

There has been a lot of speculation about whether or not various antidepressant medications are connected with greater or poorer compliance. A naturalistic analysis using claims data from 2000 patients revealed that individuals on tricyclic antidepressants may have worse compliance, and that providing family, group, or individual psychotherapy may increase compliance (Tal-Seale, Croghan and Obenchain, 2000).

Furukawa et al. did a meta-analysis of studies comparing antidepressant-benzodiazepine combos with antidepressants given alone for up to 8 weeks and found a modest benefit to co-prescribing benzodiazepines. Any potential benefit must be balanced against the clinical risks, such as the development of benzodiazepine dependency.

Depression can be effectively treated with medicine. Effective communication between doctors and patients is critical. Families and their ill relatives can benefit from education that teaches them how to manage medicine and, therefore, improve compliance, spot warning signs and symptoms, and promote an atmosphere favourable to healing and sustaining good mental health. Educational initiatives to promote compliance are founded on the assumption that poor compliance is caused by a lack of information. Psycho-educational tactics are intended to both motivate and educate patients about their disease and treatment. Because textual material alone is sometimes insufficient in long-term therapy, patients must comprehend what is expected of them. Reminder schedules, pharmacy-generated refill reminders, and customised prescription containers or packaging have all been demonstrated to greatly enhance compliance.

Counseling from a health expert might also be beneficial. Aspects of physician–patient communication that are taken into account in terms of medication adherence include: providing collaborative communication in which patients' needs are addressed, assessing patients' information and reducing the possibility of receiving conflicting information about medications; and adapting patient-centered communication to patients with mental illness in which issues about their illness and medications are discussed openly (Al-Jumah and Qureshi, 2012).

Health experts monitor and review therapy on a continuous basis. Healthcare practitioners must be aware of and concerned about the amount of patient noncompliance and its impact on treatment quality.



They should be taught about the use of medications, illness management in collaboration with patients, multidisciplinary care, and adherence monitoring. Family members must also conduct research. Family members may be a vital resource for people suffering from major mental diseases. They may help their loved ones through the diagnosis and beyond by learning more about the condition.

Psychotherapy, which is a procedure in which skilled mental health experts assist individuals in coping with their condition, is also used. This is done by talking through ways of understanding and dealing with their symptoms, thoughts, and behaviours. The most common types of psychotherapy include supportive, cognitive-behavioral, interpersonal, group, and family therapy. Certain treatments, such as art therapy or play therapy, may be beneficial, particularly for young children who have difficulty conveying their thoughts and feelings.

#### **Psychosocial factors affecting drug compliance include the following:**

**Self-Stigma:** Over the last few years, self-stigma connected with mental illness has remained a global public health problem. Patients with mental illnesses frequently have social, psychological, economic, and psychological issues as a result of self-stigma. This has an impact on, among other things, social relationships, healthcare, productivity, and acceptance (Hansson, Stjernswärd, and Svensson, 2014). It might emerge as the death of friends and loved ones—the people who are most important in one's social support network. It exacerbates mental illness and leads to social marginalisation, incapacity to participate in vital daily activities, and a low proclivity to seek treatment, ultimately affecting one main factor of quality of life (Asrat, Ayenalem, and Yimer, 2018). Self-stigma is a barrier to adequate therapy and rehabilitation for mentally ill people, resulting in longer recovery, various psychological consequences, and substantial financial issues (Adewuya et al., 2011).

**Age:** It might be difficult for health care workers to identify mental illnesses in youngsters. Children, unlike adults, go through various physical, mental, and emotional changes as they go through their normal growth and development. They are also learning to cope with, adapt to, and relate to others and the environment around them. Some problems, such as anxiety disorders, eating disorders, mood disorders, and schizophrenia, can affect both children and adults. Others begin exclusively in childhood, though they can last throughout maturity. Symptoms of obsessive-compulsive disorder, for example, occur during childhood or adolescence. It is not uncommon for a youngster to have many disorders. Age is a determinant in medication adherence since children, for obvious reasons, require more care from health professionals than adults. This, in turn, influences the rate of compliance. In a study of middle-aged and older outpatients with schizophrenia, cognitive functioning, particularly conceptualization and memory, was the

strongest patient-related predictor of medication management ability, outweighing the effects of age, gender, educational level, symptom severity, and attitudes toward medication (Jeste et al., 2003).

**Marital status:** The presence of a domestic connection with a spouse aided in health management. As a result, it was critical to evaluate the influence of marital status on medication adherence. This was due to the fact that it was seen as a measure of a social network that served as a supporting component for patients during sickness management (Sperber, Sandelowski, & Voils, 2013). Spousal support in medication compliance has been proven to give practical assistance, such as reminding the patient to take their pills (Trivedi et al., 2008).

**Economic Situation:** For individuals with mental illnesses, economic status was related to medicine compliance. The capacity to pay for the prescriptions was connected to this connection. Drugs for treating mental diseases range in price from cheap to costly (Mitchell and Selmes, 2007).

**Physician-patient Relationship:** Physician-patient relationships influence patient health outcomes such as medication adherence. Communication and continuity of care are two critical considerations. Verbal and nonverbal communication, task-oriented behaviour (effective questioning and information transmission), psychosocial behaviour (showing empathy and care), cooperation, and engagement throughout the encounter are all components of physician-patient communication processes. Aspects of physician-patient communication that are taken into account in terms of medication adherence include: providing collaborative communication in which patients' needs are addressed, assessing patients' information and reducing the possibility of receiving conflicting information about medications; and adapting patient-centered communication to patients with mental illness in which issues about their illness and medications are discussed openly (Al-Jumah and Qureshi, 2012).

**Beliefs:** According to the data, patients' beliefs had a substantial and crucial impact on affecting drug adherence. Medication usage attitudes were divided into two categories: general beliefs and specialised beliefs. Medication usage is associated with damage and overuse in general. Beliefs concerning damage included the view that pharmaceuticals were dangerous, poisonous, addictive, and should not be taken on a regular basis. Specific beliefs are those that impact patients' appraisals of the prescribed drug, such as the impression of personal need for treatment and concern about the meds' long-term use (Horne et al., 2013; Horne et al., 2001). The patients' views on the value of taking drugs mirrored their understanding of their condition and the importance of the medications in treating their sickness. As a result, their motives for taking the drug were impacted. Patients' fears about medicine, on the other hand, stemmed from



misunderstandings about what they thought they understood or had experienced with reference to the drugs.

### Hypotheses

**Ho1:** There is no significant relationship between drug compliance and the psychological wellbeing of mentally ill patients.

**Ho2:** There is no significant relationship between drug compliance and the social wellbeing of mentally ill patients.

### Methodology

The descriptive research approach was used in this study, and the population for the study included mental health patients at Adeoyo Hospital in Ibadan. However, the population's mental diagnoses were limited to schizophrenia, mood disorders, and anxiety disorders. Purposive sampling was employed as a sample strategy. The data gathering tool was a questionnaire with five components. Section A contains the demographic variable section, which

contains the demographic information about the participants that is relevant to the investigation. Section B contains remarks regarding the respondents' psychological well-being, which was measured using an eighteen-item scale adapted from Ryff's Scale of Psychological Well Being (RSPWB, 2005). Section C includes statements assessing the respondents' social well-being, which is a fifteen-item measure adapted from Lee Keyes' "Keyes Social Wellbeing Scale" (2013). Section D contains statements that assess respondents' drug compliance using a 10-item scale adapted from Hogan, Awad, and Eastwood's Drug Attitude Inventory (DAI-10) (1983). The data collection instrument (questionnaire) was verified using both face and content validity.

### Results

**Hypothesis one:** There is no significant relationship between Drug compliance and the Psychological wellbeing of mentally ill patients in Adeoyo State Hospital

**Table 1:** Pearson Product Moment Correlation (PPMC) showing the relationship between drug compliance and the psychological wellbeing of mentally ill patient

Variables	Mean	Std. Dev.	N	r	p-value	Remarks
Psychological wellbeing	48.62550	5.95228	120	.275*	.002	Sig.
Drug compliance	26.7917	4.02283				

\* Correlation is significant at the 0.05 level (2-tailed).

Table 1 showed that there is a significant relationship between drug compliance and the psychological wellbeing of mentally ill patients in Adeoyo State Hospital ( $r=.275$ ,  $n=120$ ,  $p(.002)<.05$ ). Hence, drug compliance influenced/enhanced the psychological wellbeing of mentally ill patients in the study. The hypothesis is rejected.

**Hypothesis two:** There is no significant relationship between drug compliance and social wellbeing of mentally ill patients

**Table 2:** Pearson Product Moment Correlation (PPMC) showing the relationship between drug compliance and social wellbeing of mentally ill patient

Variables	Mean	Std. Dev.	N	r	p-value	Remarks
Social wellbeing	39.4500	5.23859	120	.327*	.001	Sig.
Drug compliance	26.7917	4.02283				

\* Correlation is significant at the 0.05 level (2-tailed).

Table 2 showed that there is a significant relationship between drug compliance and social wellbeing of mentally ill patients ( $r=.327$ ,  $n=120$ ,  $p(.001)<.05$ ). Hence, drug compliance influenced/enhanced the social wellbeing of mentally ill patients in the study.

**The hypothesis is rejected.**

### Discussion of the results

Hypothesis one revealed a substantial association between medication compliance and the

psychological well-being of mentally ill individuals. This is consistent with the findings of Stentzel, van den Berg, and Schulze (2018), who conducted research to investigate possible predictors of non-adherence in patients with severe mental illnesses. According to the study's findings, the most prevalent reason for non-adherence was forgetting to take the medication. Age, on the other hand, was a detriment. Furthermore, Vaingankar, Abidin, and Chong (2020) conducted a study to determine if mental problems were connected



with present marital status. The survey was a cross-sectional nationwide survey with 6126 respondents. A systematic questionnaire was used to determine a lifetime diagnosis for five mental diseases. According to Witz (2021), males achieve professional stature and inclusion as a result of their economic, political, and prestige advantages, which they employ to limit women's access to professions. Scott and Marshall (2020) defined occupation as an economic function distinct from home activity as a result of the expansion of the labour market. Orubuloye and Oni (2018) revealed patent medicine stores where personal interactions between consumers and store owners, free consultations, and flexible pricing serve as an alternative source of treatment for many.

Hypothesis two revealed a substantial association between medication compliance and the social well-being of mentally ill individuals. The findings support Al-Jumah and Qureshi's (2012) contention that physician-patient interactions influence patient health outcomes, particularly medication adherence. Communication and continuity of care are two critical considerations. Verbal and nonverbal communication, task-oriented behaviour, psychosocial behaviour, partnership, and engagement throughout the encounter are all components of physician-patient communication processes. Aspects of physician-patient communication that are taken into account in terms of medication adherence include: providing collaborative communication in which patients' needs are addressed, assessing patients' information and reducing the possibility of receiving conflicting information about medications; and adapting patient-centered communication to patients with mental illness in which issues about their illness and medications are discussed openly. The findings are consistent with Polaris Teen's (2018) finding that mental disorders are associated with the stress of body changes, including the fluctuating hormones of puberty, as well as teen ambivalence toward increased independence and changes in their relationships with parents, peers, and others.

### Conclusion

The incidence of mental disease affects people of all ages and genders. Individuals suffering from mental illnesses have a history of noncompliance with medications and treatment. According to this study, drug compliance, on the other hand, enhances the psychosocial well-being of mentally ill patients.

### Implications for Social Casework

1. Because of the long-term usage of medications, social workers should develop an acceptable drug adherence technique for patients to follow.
2. Social workers must work with patients to change their attitudes regarding drug compliance.

3. Social workers should educate mental health personnel in hospitals about the need to use a shared decision-making care approach.
4. In order to check drug compliance, patients with mental illnesses should be observed by family members and health personnel, particularly medical social workers.
5. Medical social workers will need to educate the general public and society on the consequences of drug noncompliance.

### References

- Adewuya A. O, Owoeye A. O, Erinfolami A, Ola B. A. 2011. Correlates of Self-Stigma among Outpatients with Mental Illness in Lagos, Nigeria. *Int J Soc Psychiatry*. 57(4):418-427.
- Asrat B, Ayenalem A. E, Yimer T. 2018. Internalized Stigma among Patients with Mental Illness attending Psychiatric follow-up at Dilla University Referral Hospital, Southern Ethiopia. *Psychiatry J*. <https://doi.org/10.1155/2018/1987581>.
- Al-Jumah K. A, Qureshi N. A. 2012. Impact of Pharmacist Interventions on Patients' Adherence to Antidepressants and Patient-Reported Outcomes: A Systematic Review. *Patient Prefer Adherence*; 6: 87-100.
- Hansson L, Stjernswärd S, Svensson B. 2014. Perceived and Anticipated Discrimination in People with Mental Illness: An Interview Study. *Nord J Psychiatry*. 68(2):100-106.
- Haynes, R.B., Ackloo, E., Sahota, N., McDonald, H.P. and Yao, X. 2008. Interventions for Enhancing Medication Adherence (Review). *The Cochrane Library* (2). Wiley Publishers.
- Horne, R., Weinman, J., Barber, N., Elliot, R. & Morgan, M. 2005. Concordance, Adherence and Compliance in Medication taking: Report for the National Coordinating Centre for NHS Service Delivery and Organisation. United Kingdom: NCCSDO.
- Hunt G, Bergen J, Bashir M. 2002. Medication Compliance and Co-morbid substance abuse in schizophrenia: Impact on Community Survival 4 years after a relapse. *Schizophrenia Research* 54: 253-264.
- Janssen, B., Gaebel, W., Haerter, M., Komaharadi, F., Lindel, B. and Weinmann, S. 2006. Evaluation of Factors Influencing Medication Compliance in Inpatient Treatment of Psychotic Disorders. *Psychopharmacology*, 187 (2) 229-236.
- Jeste S. D, Patterson T. L, Palmer B. W, Dolder C. R, Goldman S, Jeste D. V. 2003. Cognitive Predictors of Medication Adherence among Middle-Aged and Older Outpatients with Schizophrenia. *Schizophr Res*. 63 (1-2): 49-58.
- Mitchell A. J, Selmes T. 2007. Why don't Patients take their Medicine? Reasons and Solutions in



- Psychiatry. *Adv Psychiatr Treat.* 2007; 13(5): 336–346.
- Mueser, K. & Gingerich, S. 2006. *The Complete Family Guide to Schizophrenia.* New York: The Guilford Press.
- Myers E. D, Branthwaite A. 1992. Out-Patient Compliance with Antidepressant Medication. *British Journal of Psychiatry*, 160:83–86.
- Noetzel, M.J., Jones, C.K. and Conn, P.J. 2012. Emerging Approaches for Treatment of Schizophrenia: Modulation of Glutamergic Signalling. *Discovery Medicine.* 14(78), 335–43.
- Novick, D., Haro J.P., Suarez, D., Perez, V., Dittann, R.W. and Haddad, P.M. 2010. Predictors and Consequences of Adherence with Antipsychotic Medication in the Outpatient Treatment of Schizophrenia. *Psychiatry Research*, 176(2–3), 109–113.
- Okunrinboye H.I., Otakpor, A.N, & Ilesanmi, O.S. 2019. Depression and medication-adherence in patients with hypertension attending a tertiary health facility in South-West Nigeria. *Pan African Medical Journal (ISSN: 1937-8688).*
- Rosa, M.A., Marcolin, M.A. and Elkis, H. 2005. Evaluation of the Factors Interfering with Drug Treatment Compliance among Brazilian Patients with Schizophrenia. *Revista Brasileira Psiquiatria*, 27(3), 178-84.
- Stentzel, U., van den Berg, N., & Schulze, L.N. 2018. Predictors of medication adherence among patients with severe psychiatric disorders: findings from the baseline assessment of a randomized controlled trial (Tecla). *BMC Psychiatry* 18, 155
- Swaminath, G. 2007. You can lead a horse to the water. *Indian Journal of Psychiatry*, 49(4), 228-230.
- Tai-Seale M., Croghan T. W, Obenchain R. 2000. Determinants of Antidepressant Treatment Compliance: Implications for Policy. *Medical Care Research and Review*, 57:491–512.
- Vaingankar, J.A., Abidin, E., & Chong, S. 2020. The association of mental disorders with perceived social support, and the role of marital status: results from a national cross-sectional survey. *Arch Public Health.*
- World Health Organization. 2002b. 25 Questions and Answers on Health and Human Rights. Health and Human Rights Publication Series, Number One. Geneva: World Health Organization.
- World Health Organization. 2014. Mental Health Policies and Programmes in the workplace. WHO Mental Health and Substance Use.